Theory of a Single Magnetic Impurity on a Thin Metal Film in Proximity to a Superconductor

Jon Ortuzar,¹ Jose Ignacio Pascual,^{1,2} F. Sebastian Bergeret,^{3,4} and Miguel A. Cazalilla^{4,2}

¹CIC nanoGUNE-BRTA, 20018 Donostia-San Sebastián, Spain

²Ikerbasque, Basque Foundation for Science, 48013 Bilbao, Spain

³Centro de Física de Materiales (CFM-MPC) Centro Mixto CSIC-UPV/EHU, E-20018 Donostia-San Sebastián, Spain

⁴Donostia International Physics Center (DIPC), 20018 Donostia-San Sebastian, Spain

(Dated: May 1, 2023)

We argue that the formation of Yu-Shiba-Rusinov excitations in proximitized thin films is largely mediated by a type of Andreev-bound state named after de Gennes and Saint-James. This is shown by studying an experimentally motivated model and computing the overlap of the wave functions of these two subgap states. We find the overlap stays close to unity even as the system moves away from weak coupling across the parity-changing quantum phase transition. Based on this observation, we introduce a single-site model of the bound state coupled to a quantum spin. The adequacy of this description is assessed by reintroducing the coupling to the continuum as a weak perturbation and studying its scaling flow using Anderson's poor man's scaling.

I. INTRODUCTION

The presence of impurities on superconductors results in subgap bound states known as Yu-Shiba-Rusinov (YSR) states¹⁻³ (see e.g. Ref. 4 for a review). These excitations can be probed using scanning tunneling spectroscopy (STS) and appear as narrow resonances in tunneling spectra.^{5,6} YSR states were originally discovered as solutions to the scattering problem of a magnetic impurity in bulk superconductors, by treating the magnetic exchange with the impurity as a classical Zeeman field that couples to the local spin density of quasi-particles.¹⁻⁴ However, this approximation does not take into account the quantum nature of the impurity spin, which can give rise to many-particle effects such as the Kondo effect⁷ and it is determinant when e.g. describing the spin carried by the YSR excitations.⁸

A fully quantum-mechanical treatment of this problem aimed at providing a comprehensive description of experiments,⁹ often requires the use of sophisticated but numerically costly methods such as the numerical renormalization group (NRG)^{10,11} or continuous time MonteCarlo.¹² In recent years, single-site models^{13–15} have emerged as a computationally affordable approach to treat some of the quantum many-particle aspects of the YSR problem.¹⁵ These models have already been successfully used to explain some spectral features observed in recent experiments.^{16–18} Moreover, it has been also applied to explain the complex many-body physics of a magnetic molecule on a clean gold film proximitized by a superconducting substrate.¹⁸

Proximitized systems have been studied mainly in the diffusive limit using the Usadel formalism.¹⁹ This approach predicts the decay of the proximity effect as well as spectroscopic features such as the closing of the gap and the formation of a minigap.^{20,21} Experiments with diffusive systems 2^{1-24} have clearly confirmed those predictions. However, the systems studied in this context are mesoscopic in size and the experimental probes that

have been employed cannot resolve the behavior of a single magnetic impurity.

On the other hand, thanks to currently available growth techniques, it is possible to grow clean metallic overlayers with thicknesses of few atomic layers on top of superconductors.^{18,25–27} These novel hybrid systems open the door to otherwise impossible on-surface synthesis, and may allow one day the study of self-organized spin chains^{28–30} as well as other, more complex, molecular structures³¹ on superconductors. Such systems are



Figure 1. Panel (a) shows a sketch of the studied system: a magnetic impurity interacting via exchange with a proximitized thin metal film. Panels (b) and (c) are the local density of states (LDoS) in the bulk of the superconductor and in the thin proximitized metallic film. Panel (d) is a convoluted⁶ spectroscopic measurement using STM of a thin (about 4 mono-layers) proximitized Au film on V(100)

clearly not in the diffusive limit and have to be described within the ballistic limit. In this case, subgap bound states appear in the normal region and extend into the superconductor over distances of the order of the coherence length. The existece of such states has been known for some time, since the work of de Gennes and Saint James.^{32,33}

The aim of this work is to extend the application of the single-site model¹⁵ to describe the complexity of a single magnetic impurities interacting with a thin metallic film in proximity to a superconductor. This is a problem of much interest to a number of recent experiments.^{18,34,35} Below, we first study the system treating the magnetic impurity as a classical spin in the ballistic limit where there is a single de Gennes-Saint James (dGSJ) bound state in the gap. We find that a large overlap exists between the wavefunctions of the dGSJ and YSR states. Motivated by this result, we propose that the single-site model is a relevant simplified model for complex system consisting of the magnetic impurity on the proximitized thin film. The model can be solved exactly and also provides a computationally cheap way to treat the manyparticle effects associated with the quantum spin of the impurity. The adequacy of the single-site model for the system of interest here is assessed by means of a "poor man's" scaling analysis. To this end, we introduce a Hamiltonian consisting of a single-site model perturbed by an impurity-mediated coupling to the continuum of other excitations. Under certain conditions we find that, as the high-energy continuum states are integrated out, the impurity remains most strongly coupled to the single site describing the dGSJ state.

The structure of the article is the following: In Sec. II we describe the system and the approximations used. Section III is divided in two subsections with the first focusing on the YSR states resulting from the interaction of the magnetic impurity with the proximitized film. This study is undertaken assuming the spin of the magnetic impurity can be treated classically. In the second subsection, we describe the calculation of the wave function overlap between the YSR and dGSJ states. In Sec. IV, we introduce the single-site model for the magnetic impurity on a proximitized film. Finally, in section V, we argue that the single-site model provides an accurate description of this system using Poor man's scaling.³⁶ The most technical details of the calculations have been relegated to the Appendices.

II. SYSTEM AND MODEL

Fig. 1 shows a schematic picture of the system studied in this work, which is motivated by experiments reported in Ref. 18 and Refs. 25–27. The system consists of a magnetic impurity on top of a thin normal metal film (N) in proximity to a superconductor (S). The superconductor occupies the half-space x > 0, while the N film corresponds to -a < x < 0. The system is translationally invariant in the (y, z)-plane, so it is convenient to describe the electron wave function as $\psi(x, \mathbf{k}_{\parallel})$, where \mathbf{k}_{\parallel} is the component of the momentum vector parallel to the S/N interface at x = 0. We assume a perfect S/N interface with no Fermi surface mismatch or potential barrier, such that the N region acts as a cavity for electrons with energy $E < \Delta$: they undergo Andreev retro-reflections at the S/N interface and normal specular reflections at the interface with vacuum. According to the Bohr-Sommerfeld quantization rule, the phase accumulated along a closed classical trajectory must be a multiple of 2π . In the N/S system under consideration, a closed trajectory consists of two Andreev retroreflections at the S/N interface and two normal reflections at x = -a. Thus,

$$\frac{2aE}{\hbar v_F \cos\varphi} - \cos^{-1}\left(\frac{E}{\Delta}\right) = n\pi \tag{1}$$

where $\cos^{-1}(E/\Delta)$ is the phase shift associated to each AR, and $\cos \varphi = k_{\parallel}/k_F$. Eq. (1), determines the subgap bound states, also known as De Gennes-Saint James (dGSJ) states.³² It is valid for clean N-layers with a mean free path larger than the thickness *a*, and it describes a continuum of subgap states.^{32,37}

For the STM experiments of interest to us here, assuming specular tunneling,³⁸ the decay of the wavefunction of these excitations in vacuum is determined by the metal work-function. This energy scale is of the order of one electron-volt and therefore much larger than the superconducting gap. Therefore, in vacuum, the tail of the dGSJ wave function is essentially indistinguishable from that of an electron at the Fermi level in the normal state, and excitations with finite k_{\parallel} penetrate less into the vacuum. As a result, when probed with a STM in the tunneling regime, excitations with large $|\mathbf{k}_{\parallel}|$ are filtered out^{38–40} and dGSJ states are observed as narrow subgap resonances made of dGSJ quasi-particles with $k_{\parallel} \approx 0.^{18}$ Moreover, a small amount of disorder will randomize trajectories with $\cos \varphi < a/l$, where l is the mean free path, suppressing the coherence of such trajectories. A magnetic impurity on top of the proximitized film has compact and anisotropic orbitals that typically couple to several scattering channels from the substrate. However, since the dGSJ quasi-particles with $k_{\parallel} \simeq 0$ penetrate farther into the vacuum, they are also expected to contribute substantially to the most strongly coupled scattering channel. Thus, one can effectively approximate the tunneling problem using a one-dimensional model which neglects the motion parallel to the surface:

$$\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_J , \qquad (2)$$

where

$$\mathcal{H}_{0} = \sum_{\sigma} \int_{-a}^{\infty} dx \,\psi_{\sigma}^{\dagger}(x) \left[-\frac{\hbar^{2}}{2m^{*}} \partial_{x}^{2} - E_{F} \right] \psi_{\sigma}(x') + \int_{0}^{\infty} dx \,\Delta \,\psi_{\uparrow}^{\dagger}(x) \psi_{\downarrow}(x) + \text{h.c.} , \qquad (3)$$

and

$$\mathcal{H}_J = \sum_{\sigma\sigma'} J \psi_{0\sigma}^{\dagger} \boldsymbol{S} \cdot \boldsymbol{s}_{\sigma\sigma'} \psi_{0\sigma'} .$$
 (4)

Here, $\psi_{\sigma}(x)$ ($\psi_{\sigma}^{\dagger}(x)$) represents the annihilation (creation) operator for an electron with spin $\sigma = \uparrow, \downarrow$ in the metal-superconductor substrate. H_0 describes a proximitized thin film of thickness a > 0. The first term contains the kinetic energy and chemical potential E_F , and the second term is the s-wave pairing potential. The pairing potential is not self-consistently calculated. Corrections due to self-consistency result in a spatially non-uniform pairing potential $\Delta(x)$, but they have only a small effect on the spectral properties of the dGSJ states.^{40,41} The magnetic exchange with the impurity is described by H_J , with s denoting the electron-spin Pauli matrices and S denoting the impurity spin operator. The operators $\psi_{0\sigma}$ ($\psi_{0\sigma}^{\dagger}$) annihilate (create) electrons at the position of the impurity. For the one-dimensional model introduced above, $\psi_{0\sigma} = \psi_{\sigma}(x = -a)$. In the following section, we analyze this model using the approach of Yu, Shiba, and Rusinov (YSR),^{1–3} where the impurity spin \boldsymbol{S} is treated as a classical vector.

III. YSR IN PROXIMITIZED THIN FILMS

In the previous section, we have derived the equation that determines the spectrum of subgap states (cf Eq. 1) using the Bohr-Sommerfeld semiclassical approximation. As explained above, we will focus on the one-dimensional case, which corresponds to $\cos \varphi = 1$ in Eq. (1). To deal the coupling to the magnetic impurity, we solve the model described by Eqs. (2)-(4). To this end, we use Green's functions (GFs) and follow the approach outlined in Ref. 40. The technical details of the calculation are described in Appendix A. From the knowledge of the retarded GFs, $G(\omega + i\eta, x, x)$, the local density of states (LDoS) $\rho(\omega, x)$ of the system is obtained by using $\rho(\omega, x) = -\frac{1}{\pi} \operatorname{Im} G(\omega + i\eta, x, x)$.

Fig. 2 (a) shows the LDoS on the surface as a function of film thickness. As we increase the thickness, new dGSJ states enter the gap. The GF also has poles with a finite imaginary part outside the superconducting gap that correspond to states in the continuum (i.e. above the superconductor gap), and give rise to McMillan-Rowell-Tomasch oscillations.^{42,43} From here on, we focus our discussion on thin films with a single subgap bound state. In the following subsection, we tackle the coupling to the magnetic impurity.

A. YSR states

The GF for the S/N system provides the starting point for calculating the spectral properties of the YSR excitations. The properties of the latter can be obtained by solving the following integral equation:

$$G_{YSR}(x, x') = G(x, x') + G(x, -a)V \times (1 - VG(-a, -a))^{-1}G(-a, x').$$
(5)

Here, G(x, x') is the GF obtained in the previous subsection. The scattering potential for a spin-S impurity in the Nambu notation is $V = JS\sigma_z\tau_0$, assuming that the impurity (classical) spin points along the z-axis.

Fig 2 (b) shows the evolution of the YSR state as a function of the exchange coupling $\alpha = \nu_0 \pi J S$, with ν_0 being the normal metal DoS defined so that the quantum phase transition (QPT), where the energy of the YSR state crosses the center of the gap, happens for $\alpha = 1.^{1-3}$ Note that the exchange coupling splits dGSJ state into two states (spin up and down), one of which shifts to higher energy while the other shifts to lower energy, see Fig. 2(c). As J increases beyond a certain value, the higher energy state disappears into the continuum. From this point on, the energy of the remaining subgap state behaves similarly to a YSR in a bulk superconductor.^{1–3} For thicker films, with more than one dGSJ state, the behavior is similar: each bound state splits in two, shifting in opposite directions depending on their spin projection, with more excited states eventually merging in the continuum and disappearing.

The transmutation of the dGSJ into the YSR state can be regarded as a consequence of a spectral reorganization taking place around $|\omega| = \Delta$ caused by AR (see Fig. 1c). Using an analogy to semiconductor physics, YSR states appear in a superconductor because the coherence "peak" behavior ~ $(\omega^2 - \Delta^2)^{-1/2}$ (cf. Fig.1 b) resembles a van Hove singularity at the bottom (top) of the conduction (valence) band of a one-dimensional insulator. Bound states appear due to the infinitesimal attraction provided by the magnetic impurity Dirac-delta potential. However, in a proximitized film, AR reorganizes the spectral weight by removing the van Hove-like singularity while shifting most of its spectral weight to the dGSJ state (cf. Fig.1 c). Together with the localization of the dGSJ states at the surface, this enables the transmutation of one of the dGSJ states per spin into a YSR. Thus, a large overlap of the wavefunctions of YSR and dGSJ states is expected, as explicitly demonstrated in the following subsection.

B. Overlap between SJdG and YSR States

In this section, we compute the overlap of the YSR and the dGSJ states as a function of the exchange coupling J. This can be achieved by using the GF obtained from the scattering solution of the problem with and without



Figure 2. (a) Evolution of the dGSJ states as a function of the thickness of the metallic layer. (b) Evolution of the YSR state for a fixed metallic layer thickness as a function of the exchange coupling. (c) Zoom-in of the evolution of the YSR state. (d) Wavefunction of the SJdG and YSR states averaged over distances $\gg k_F^{-1}$.

magnetic impurity. The square of the overlap is computed from the following integral involving the residue of the two GFs:

$$\Theta|^{2} = \int dx \left[u_{dGSJ}(x) u_{YSR}^{*}(x) + v_{dGSJ}(x) v_{YSR}^{*}(x) \right]$$
$$= \int dx dx' \operatorname{Tr} \left\{ \operatorname{Res} \, G(x, x') \right) \operatorname{Res} \, G_{YSR}(x', x) \right\}.$$
(6)

Here $\operatorname{Res} G_{YRS}$ ($\operatorname{Res} G$) is the residue of the Nambu GF matrix at the YSR (dGSJ) pole with spin up.

In Fig. 3 (a) and (b) we show the behavior of the overlap Θ as a function of exchange coupling J for different values of film thickness (which determines the dGSJ state energy). To check our results beyond the leading order in Δ/E_F , we also compute the overlap by solving the Bogoliubov-de Gennes equations for a one-dimensional tight-biding chain containing up to 2000 sites. The results are shown in Fig. 3(c) as a function of J normalized to the critical value J_c where the system undergoes the parity-changing quantum phase transition.¹⁻⁴ The overlap between the YSR and dGSJ states decreases as the exchange coupling increases, but it remains close to unity even across the quantum phase transition. It is worth noting that the energy of the YSR excitation shifts away from that of the dGSJ state as the exchange coupling is increased. The significant overlap between the two states suggests that the YSR state primarily descends from the dGSJ state, with a minor contribution from the continuum states of the proximitized film. Therefore, in a first approximation, the coupling with the magnetic impurity can be described by replacing the proximitized film with a single level representing the dGSJ state.

IV. SINGLE-SITE MODEL

Motivated by the results of the previous section, we introduce a simplified model that replaces the entire proximitized film with a single site representing the dGSJ state. As we show below, this model is useful for analyzing the coupling between the dGSJ state and a quantum spin. The Hamiltonian of the single site is given by:

$$H_0 = \sum_{\sigma} E_s \left(\gamma_{\sigma}^{\dagger} \gamma_{\sigma} - \frac{1}{2} \right), \tag{7}$$

where γ_{σ} ($\gamma_{\sigma}^{\dagger}$) are the annihilation (creation) operators for a dGSJ quasi-particle with spin $\sigma = \uparrow, \downarrow$, and E_s is the eigenvalue of the BdG Hamiltonian (in the absence of magnetic impurity). As explained in Appendix B, this Hamiltonian can be recast in terms of electron operators $d_{\sigma}, d_{\sigma}^{\dagger}$ as follows:

$$H_0 = U \sum_{\sigma} n_{\sigma} + \left[\Delta_s d_{\downarrow} d_{\uparrow} + \text{h.c.} \right], \qquad (8)$$

where $n_{\sigma} = d_{\sigma}^{\dagger} d_{\sigma}$; U and Δ_s are effective scattering and pairing potentials, respectively. In terms of U and Δ_s , $E_s = \sqrt{U^2 + \Delta_s^2}$. Without loss of generality, below we discuss the particle-hole symmetric case where U = 0 and therefore $E_s = \Delta_s$

Next, we introduce the coupling to the impurity. To make contact with the classical description employed in the previous section, we first discuss the Ising limit of the exchange coupling, i.e.

$$H_J^{\text{Ising}} = J_{dd}^{\parallel} S^z \left(n_{\uparrow} - n_{\downarrow} \right), \qquad (9)$$

where $J_{dd}^{\parallel} > 0$ is the exchange coupling with the dGSJ quasi-particle. This model reproduces the most salient features of the YSR states described above. To begin with, note that, besides the fermion parity $P = \prod_{\sigma} (-1)^{n_{\sigma}} = \pm 1$, the impurity spin operator S^z is also conserved in this limit, i.e. $\left[S^z, H_0 + H_J^{\text{Ising}}\right] = 0$. Thus, the ground state is doubly degenerate corresponding to the two possible orientations of the classical vector $\mathbf{S} = \pm S \hat{\mathbf{z}}$: For $J_{dd} < J_c = 2\Delta_s$ the ground state is one of the two following states $\{|BCS\rangle \otimes |\pm \frac{1}{2}\rangle\}$ with P = +1



Figure 3. Overlap between the YSR wavefunction and the SJdG wavefunction. Panels (a) and (b) show the calculation done from the continuous model for different SJdG bound state energies and values of E_F/Δ . Panel (c) shows the same calculation done with a tight-binding model.

and $\gamma_{\sigma}|BCS\rangle = 0$. For $J_{dd} > J_c$, the ground state is one in $\{|\uparrow\rangle \otimes |-\frac{1}{2}\rangle, |\downarrow\rangle \otimes |+\frac{1}{2}\rangle\}$ with P = -1 and $|\sigma\rangle = \gamma_{\sigma}^{\dagger}|BCS\rangle$. The YSR excitation is a transition between these two ground states of opposite parity with excitation energy^{15,18} $|\Delta_s - J_{dd}/2|$. In addition, the odd parity sector of the Hilbert space also contains the following two states: $\{|\uparrow\rangle \otimes |+\frac{1}{2}\rangle, |\downarrow\rangle \otimes |-\frac{1}{2}\rangle\}$ with excitation energy equal to $\Delta_s + J_{dd}/2$. For small J_{dd} , a transition from the ground state with P = +1 to these states corresponds to the second subgap peak in the LDoS of the classical approach that shifts up in energy with increasing exchange and eventually disappears into the continuum, see Fig 2 (c).

Next, we generalize Eq. (9) by adding the spin-flip term, which allows the impurity spin to fluctuate:

$$H_J^d = J_{dd}^{\parallel} S^z \left(n_{\uparrow} - n_{\downarrow} \right) + J_{dd}^{\perp} (S^+ d_{\downarrow}^{\dagger} d_{\uparrow} + \text{h.c.})$$
(10)

As argued in Refs. 14 and 15, the single-site model provides an economical and fully quantum-mechanical description of YSR spectra in superconductors which compares well with the results obtained using sophisticated but computationally expensive methods like the Numerical Renormalization Group (NRG).¹¹ The accuracy of this description in the present system will be addressed in the following section.

The spin-flip term, $\propto J_{dd}^{\perp} > 0$, has important consequences for the spectrum of the model. In the weak coupling limit, i.e. for $J_{dd}^{\parallel} + 2J_{dd}^{\perp} < 2\Delta_s$, (assuming an unbiased preparation of the system) the ground state is described by the following density matrix:

$$\rho_{GS} = \frac{1}{2} \left[|BCS, +\frac{1}{2}\rangle \langle \langle +\frac{1}{2}, BCS| + |BCS, -\frac{1}{2}\rangle \langle -\frac{1}{2}, BCS| \right], \tag{11}$$

On the other hand, in the strong coupling limit where $J_{dd}^{\parallel} + 2J_{dd}^{\perp} > 2\Delta_s$, the ground state is a singlet:

$$|GS\rangle = \frac{1}{\sqrt{2}} \left(|\uparrow\rangle \otimes |+\frac{1}{2}\rangle + |\downarrow\rangle \otimes |-\frac{1}{2}\rangle\right).$$
(12)

that is, a pure state resulting from the quantum superposition of the two ground states of the Ising limit of the model. In weak and strong-coupling regimes, unlike the conventional classical approach of YSR,¹⁻³ the quantum model predicts that YSR excitations carry no spin polarization.⁸

Finally, since in the original model (cf. Eq. 4) the energy of the YSR does not grow without bound as the exchange with the magnetic impurity J becomes arbitrarily large, the couplings $J_{dd}^{\perp}, J_{dd}^{\parallel}$ cannot be much larger than Δ_s in the single-site model. Note that, for large $J_{dd}^{\perp}, J_{dd}^{\parallel}$

the energy of the YSR grows like $\max\{J_{dd}^{\perp}, J_{dd}^{\parallel}\}$. Thus, for the energy of the YSR to remain within the gap, the exchange couplings of the single-site model must saturate to an upper bound so that $\max\{J_{dd}^{\perp}, J_{dd}^{\parallel}\} \lesssim \Delta_s$. Therefore, they must be regarded as renormalized exchange interactions, which are also the result of the spectral reorganization and localization of excitations with energy $\sim \Delta_s$ caused by Andreev reflection at the S/N interface.

V. SCALING APPROACH

In order to investigate the accuracy of the single-site model, we reintroduce the coupling to the continuum of excitations as a perturbation. Whether this perturbation changes the low-energy spectrum substantially or not can be assessed using the poor man's scaling method,³⁶ as we

describe in the following.

In the single-site model, the effective exchange coupling of the impurity and dGSJ quasi-particle is $J_{dd} = J_{dd}^{\perp} = J_{dd}^{\parallel}$, where, for the sake of simplicity, we assume an isotropic coupling. Our conclusions also apply to the anisotropic case with small modifications. Through the exchange interaction with the magnetic impurity, the dGSJ quasi-particles can also couple to the continuum of excitations of the proximitized film. Let us introduce the following modified exchange coupling which, besides the coupling to the dGSJ, describes an impurity-mediated coupling of the dGSJ-site to the continuum, and will be treated below as a perturbation:

$$H_{J}^{dc} = \sum_{\sigma\sigma'} \left(J_{dd} \, d_{\sigma}^{\dagger} \boldsymbol{s}_{\sigma\sigma'} d_{\sigma} + J_{\Phi\Phi} \, \Phi_{0\sigma}^{\dagger} \boldsymbol{s}_{\sigma\sigma'} \Phi_{0\sigma'} \right) \cdot \boldsymbol{S} + J_{d\Phi} \sum_{\sigma\sigma'} \left(d_{\sigma}^{\dagger} \boldsymbol{s}_{\sigma\sigma'} \Phi_{0\sigma'} + \Phi_{0\sigma}^{\dagger} \boldsymbol{s}_{\sigma\sigma'} d_{\sigma'} \right) \cdot \boldsymbol{S}.$$

$$(13)$$

The operators $\Phi_{0\sigma}$, $\Phi_{0\sigma}^{\dagger}$ are the annihilation and creation operators for electrons in the continuum at the position of magnetic impurity. Phenomenologically, we have assumed different couplings for the various processes involving the scattering of the dGSJ and the continuum excitations by the impurity. These couplings can be calculated from first principles. However, they depend on microscopic details of the matrix elements of the impurity orbitals and the continuum of both subgap and outer-gap excitations which are difficult to model. For this reason, we treat their bare values as free parameters in the analysis below.

We carry out the poor man's scaling analysis³⁶ of the model (13) by integrating out the high energy degrees of freedom from the continuum with energies of the order of the bandwidth $D \sim E_F$. Since these band-edge modes exhibit vanishing superconducting correlations because their energies are well above the gap, the calculations do not differ much from those of the standard Kondo scaling of a magnetic impurity.³⁶ Some details are provided in Appendix C. In what follows, we focus on the discussion of the solutions to the scaling equations, which read

$$\frac{dg_{\Phi\Phi}}{d\ell} = g_{\Phi\Phi}^2, \qquad (14)$$

$$\frac{dg_{d\Phi}}{d\ell} = g_{d\Phi}g_{\Phi\Phi},\tag{15}$$

$$\frac{dg_{dd}}{d\ell} = g_{d\Phi}^2. \tag{16}$$

Here $g_{dd} = 2\nu_0 J_{dd}$, $g_{d\Phi} = 2\nu_0 J_{d\Phi}$, and $g_{\Phi\Phi} = 2\nu_0 J_{\Phi\Phi}$ are dimensionless couplings, $\nu_0 \sim 1/D$ being the mean density of continuum states. The scaling variable ℓ is defined such that the bandwidth is reduced according to $D(\ell) = De^{-\ell} \to 0$ as $\ell \to +\infty$, where $D \sim E_F$.

As the bandwidth of the system is reduced, the above scaling equations imply that the renormalization of g_{dd} and $g_{d\Phi}$ is driven by the growth of $g_{\Phi\Phi}$. Indeed, Eq. (14)

for $g_{\Phi\Phi}$ is mathematically identical to the scaling equation for the exchange coupling of a magnetic impurity in a normal metal (Kondo scaling). It can be readily solved by the ansatz $g_{\Phi\Phi}(\ell) = (\ell^* - \ell)^{-1}$, where $\ell^* = 1/g_{\Phi\Phi}(0)$. Like the ordinary Kondo scaling, ℓ^* corresponds to the logarithmic scale where $g_{\Phi\Phi}(\ell)$ diverges and the perturbative renormalization breaks down. This happens when the bandwidth becomes of the order of a "Kondo temperature", T_K^{Φ} , i.e. for $\ell^* = \log(D/T_K^{\Phi})$. Hence, $g_{\Phi\Phi}(\ell^*) \sim 1$ leads to $T_K^{\Phi} = De^{1/(2\nu_0 J_{\Phi\Phi})}$. Note that $T_K^{\Phi} \gg \Delta_s$ would imply that the continuum states at energies much higher than the superconducting gap are strongly coupled to the magnetic impurity. In this situation, the single-site description as introduced above breaks down. In the classical approach, such a strong coupling to the continuum should result in substantial suppression of the overlap between the YSR and dSGJ states.

Indeed, the wavefunction overlap Θ (cf. Fig. 3) can be used to obtain a rough estimate the ratios of the bare couplings $g_{d\Phi}(0)/g_{dd}(0)$, and $g_{\Phi\Phi}(0)/g_{dd}(0)$. To this end, we first notice that $g_{dd} \sim J_{dd}, g_{d\Phi} \sim J_{d\Phi}$, and $g_{\Phi\Phi}(0) \sim J_{\Phi\Phi}$ contain matrix elements with zero, one, and two powers of the continuum orbitals, respectively (recall that the exchange couplings are second order in the matrix element describing the tunneling between the impurity magnetic orbital and the metallic host states). Let $\gamma = 1 - |\Theta|$ measure the degree of admixture of the YSR state with the continuum; γ will be enhanced by quantum fluctuations relative to the estimates provided by the classical approach (cf. Sec. III). Nonetheless, we expect γ to remain much smaller than one. Thus, $g_{dd}(0) \sim \gamma^0$, $g_{d\Phi} \sim \gamma$ and $g_{\Phi\Phi} \sim \gamma^2$, to leading order in γ . Furthermore, $g_{dd}(0) = 2\nu_0 J_{dd} \sim \Delta_s / D \sim \Delta / D \ll 1$ according to the discussion at the end of the previous section.

Next, we proceed to obtain solutions to the scaling equations using the above estimates for the initial conditions of the flow. Concerning the solutions of (15) and (16), we notice that (15) is solved by the ansatz $g_{d\Phi}(\ell) = r_{d\Phi}/(\ell^* - \ell)$ with $r_{d\Phi} = g_{d\Phi}(0)/g_{\Phi\Phi}(0)$. Introducing this result into Eq. (14) and integrating, we obtain the following renormalized coupling between the impurity and the dSGJ:

$$g_{dd}(\ell) = g_{dd}(0) + \frac{g_{d\Phi}^2(0)}{g_{\Phi\Phi}(0)} \frac{(\ell/\ell^*)}{1 - (\ell/\ell^*)}$$
(17)

Using $g_{d\Phi}^2(0)/g_{\Phi\Phi}(0) = \gamma^2 g_{dd}^2(0)/[\gamma^2 g_{dd}(0)] \simeq \gamma^0 g_{dd}(0)$, the above expression simplifies to:

$$g_{dd}(\ell) \simeq \frac{g_{dd}(0)}{1 - (\ell/\ell^*)}.$$
 (18)

which needs to be compared with the behavior of the renormalized coupling to the continuum after setting $g_{\Phi\Phi}(0) \simeq \gamma^2 g_{dd}(0)$:

$$g_{\Phi\Phi}(\ell) \simeq \frac{\gamma^2 g_{dd}(0)}{1 - (\ell/\ell^*)}.$$
 (19)

Note that both couplings diverge at $\ell^* = \log(D/T_K^{\Phi})$ with $T_K^{\Phi} \simeq De^{-1/2(\nu_0 \gamma^2 J_{dd})} \ll \Delta$ if $\gamma \ll 1$, which is consistent with what was discussed above. For instance, if we choose $\gamma \approx 0.2$ (corresponding to $\Theta \approx 0.8$), then

$$\frac{g_{\Phi\Phi}(\ell)}{g_{dd}(\ell)} \simeq \gamma^2 \ll 1.$$
(20)

Thus, as the continuum states are integrated out, the impurity remains most strongly coupled to the single site describing the dGSJ quasi-particle and therefore the single-site model remains an accurate description of the magnetic impurity on the proximitized thin film.

Let us close this section by pointing out some potential problems with the scaling analysis described above. First of all, like the original poor man's scaling,³⁶ the equations are obtained perturbatively. Therefore, the solutions to the scaling equations are valid provided the couplings remain small compared to unity. This is not a problem under the above assumptions because the scale where the couplings diverge ℓ^* is much smaller than the superconductor gap and the scaling must be stopped at the scale of Δ . As we get closer to the gap scale, the superconducting correlations cannot be neglected, and taking them into account will modify the flows of the renormalized couplings. Nevertheless, we should interpret the above analysis as providing information on the tendency of the high-energy continuum states to couple to the impurity in the presence of the coupling to the dGSJ state. In order to follow the renormalization of the coupling to the continuum from high to low energies, it would be desirable to carry out calculations using the NRG and starting from a more microscopic description of the system, e.g. using model parameters obtained from first principle calculations. Such calculation should provide a more quantitative assessment of the accuracy the single-site model introduced in this work for proximitized films.

VI. CONCLUSIONS

We have studied the YSR excitations in a thin metal film proximitized by a superconductor. This has been carried out by introducing a one-dimensional model of the metal film/superconductor substrate. We have discussed the spectrum of this model, which consists of subgap bound states known as de Gennes-Saint James (dGSJ) states. We have shown that Andreeev-reflection at the metal/superconductor interface leads to a substantial spectral reorganization around and below the gap energy. Next, the spectrum of the system when a magnetic impurity is deposited on the metal film has been also described. Treating the impurity spin as a classical vector, we have found there is substantial overlap of the wavefunctions of the Yu-Shiba-Rusinov (YSR) and the dGSJ states. Motivated by these results, a singlesite model has been introduced. This model replaces the

complexity of the proximitized film with a single-site that represents the dGSJ quasi-particle excitation and is coupled to the impurity with an effective change coupling. The single-site model is exactly solvable and allows us to go beyond the classical description of the impurity by treating its spin quantum mechanically. Finally, we have addressed the accuracy of the single-site model by phenomenologically re-introducing the coupling to the continuum of excitations of the proximitized film as a perturbation and using the poor man's scaling method: Under conditions suggested by the findings of the classical approach, we have shown that the exchange coupling with the site that describes the dGSJ quasi-particle excitation remains the dominant coupling under scaling. Thus, the continuum of excitations of the proximitized film can be neglected in a first approximation, and the YSR states can be regarded as resulting from the exchange interaction of the magnetic (quantum) impurity with the dGSJ quasi-particles.

The approach used here can be generalized to treat impurities with higher spin and account for single-ion as well as magnetic exchange anisotropies. Our results provide theoretical support for the model used to analyze the STS spectra reported in Ref. 18. In addition, since the single-site model introduced here is computationally cheaper than more sophisticated numerical methods like the numerical renormalization group $(NRG)^{11}$ or continuous-time Montecarlo,¹² it can be used to model more complex systems such as chains or other nanostructures of magnetic impurities on proximitized films, which would be otherwise rather intractable by those methods. For this reason, we also believe it is worth revisiting the system studied here using much more sophisticated numerical tools, in order to quantitatively assess the limitations of the single-site model as introduced in this work.

ACKNOWLEDGMENTS

We acknowledge financial support from Grants No. PID2019-107338RB-C61, No. CEX2020-001038-M, No. PID2020-112811GB-I00, and PID2020-114252GB-I00, funded by MCIN/AEI/ 10.13039/501100011033, from the Diputación Foral de Guipuzcoa, the ELKARTEK project BRTA QUANTUM (no. KK-2022/00041), and from the European Union (EU) through the Horizon 2020 FET-Open projects SPRING (No. 863098) and SU-PERTED (No. 800923), and the European Regional Development Fund (ERDF). M.A.C. has been supported by Ikerbasque, Basque Foundation for Science, and MCIN Grant No. PID2020-120614GB-I00 (ENACT). F.S.B. thanks Prof. Björn Trauzettel for his hospitality at Würzburg University, and the A. v. Humboldt Foundation for financial support. J.O. acknowledges the scholarship PRE_2022_2_00950 from the Basque Government. The authors also thank Katerina Vaxevani and Stefano Trivini for their help measuring the experimental spectrum on Fig 1 (d) and several discussions.

Appendix A: Two-layer Green's Functions

As discussed in Ref. 44, the GFs of a composite system (3) can be obtained from the GFs of the constituent subsystems. We denote the GFs of each subsystem as $g_i(x, x')$, with i = N, S. Next, we impose the following boundary conditions:

$$\frac{dg_i(x,x')}{dx}\Big|_{x=0} = 0, \qquad \frac{dg_i(x,x')}{dx'}\Big|_{x'=0} = 0, \tag{A1}$$

$$\frac{dg_N(x,x')}{dx}\Big|_{x=-a} = 0, \qquad \frac{dg_N(x,x')}{dx'}\Big|_{x'=-a} = 0, \tag{A2}$$

$$g_S(x \to +\infty, x') = 0, \qquad g_S(x, x' \to +\infty) = 0, \tag{A3}$$

$$\lim_{\delta \to 0^+} \tau_3 \frac{dg_i(x, x')}{dx} \Big|_{x=x'-\delta}^{x=x'+\delta} = \frac{2m}{\hbar^2}, \qquad \lim_{\delta \to 0^+} \tau_3 \frac{dg_i(x, x')}{dx'} \Big|_{x'=x-\delta}^{x'=x+\delta} = \frac{2m}{\hbar^2}.$$
 (A4)

We assume a zero derivative at the vacuum interface, except for the semi-infinite superconductor at $x, x' \to \infty$, for which the GFs are assumed to vanish.

Using the above boundary conditions, and assuming continuity of the full GF and its derivative at the N/S interface, we obtain the following relations:

$$G(x,x') = \begin{cases} g_S(x,x')\theta(x') \mp g_S(x,0)[g_S(0,0) + g_N(0,0)]^{-1}g_{S,N}(0,x') , \text{ for } x > 0, x' \ge 0\\ g_N(x,x')\theta(-x') \mp g_N(x,0)[g_S(0,0) + g_N(0,0)]^{-1}g_{S,N}(0,x') , \text{ for } x < 0, x \ge 0 \end{cases}$$
(A5)

The GFs for the isolated system are easy to calculate from Bogoliubov-de Gennes equation.³⁸ After that, one can get the dressed GF from (A5), see [Arnold]. The GF shows a pole with the following energy distribution:

$$\frac{\omega}{\sqrt{\Delta^2 - \omega^2}} \tan\left(\frac{2ma}{\hbar^2 k_F}\omega\right) = 1 \tag{A6}$$

Note that by expanding the tangent around zero to the first order we arrive at the same solution as obtained from a semi-classical argument Eq. (1).

Appendix B: Single site Hamiltonian for a proximitized superconductor

By solving the Bogoliubov-de Gennes equations for the proximitized thin film, the electron field operator at the position of the magnetic impurity, $\phi_{0\sigma}$, can be written as follows:

$$\psi_{0\sigma} = u_0 \gamma_\sigma + \sigma v_0^* \gamma_{-\sigma}^\dagger + \Phi_{o\sigma}. \tag{B1}$$

We start by changing the basis on the unperturbed Hamiltonian (3), where first two terms described the dGSJ quasiparticle and $\Phi_{0\sigma}$ describes the modes in the continuum. We now introduce a rotation for the operators creating the discrete state:

$$\begin{pmatrix} d_{\sigma} \\ d_{-\sigma}^{\dagger} \end{pmatrix} = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} \gamma_{0\sigma} \\ \gamma_{0-\sigma}^{\dagger} \end{pmatrix} .$$
(B2)

This rotation leads to (8), where $U = E_s \cos 2\theta$ and $\Delta_s = E_s \sin 2\theta$. Furthermore, requiring that

$$\psi_{\uparrow} = \sqrt{Z}d_{\uparrow} + +\Phi_{0\uparrow} = u_0\gamma_{\sigma} + \sigma v_0^*\gamma_{-\sigma}^{\dagger} + \Phi_{0\uparrow} .$$
(B3)

Hence, $\tan \theta = -v_0/u_0$ and $Z = u_0^2 + v_0^2$, where $U = E_s(u_0^2 - v_0^2)$ and $\Delta_s = 2E_s u_0 v_0$.

Appendix C: Calculation of the scaling equations

In order to perturbatively obtain the scaling equations of the model introduced in Sec. V, we consider an expansion of the partition function of the system, i.e.

$$Z(D) = Z_0(D) \left\langle \mathcal{T} \exp\left[-\int_0^\beta H_J^{dc}(\tau)\right] \right\rangle_0,\tag{C1}$$

in powers of the couplings J_{dd} , $J_{d\Phi}$ and $J_{\Phi\Phi}$. In the above expression for $Z(D) Z_0(D) = \text{Tr}e^{\beta H_0}$ is the partition function of the system without magnetic impurity at inverse absolute temperature $\beta = (k_B T)^{-1}$; $\langle \dots \rangle_0$ is the expectation value over the non-interacting grand canonical ensemble defined by H_0 . The operator $H_J(\tau) = e^{H_0\tau}H_Je^{-H_0\tau}$, where H_J is given in Eq. (13), describes the magnetic exchange with the impurity in the interaction representation and \mathcal{T} is the imaginary time-ordering symbol. We have also introduced a parameter, $D E_F$, which is the bandwidth of the composite thin film and superconductor system.

Following Anderson,³⁶ we shall use perturbation theory to obtain a map onto a system with smaller bandwidth $D' = D - \delta D < D$. Associated with the bandwidths D and D', there are also the following characteristic (imaginary) time scale (in units where $\hbar = 1$) $\tau_c = D^{-1}$ and $\tau'_c = (D')^{-1} > \tau_c$. The lowest order terms of the perturbation series for the system with bandwidth D read:

$$Z(D) = Z_0(D) \left\{ 1 - \int d\tau \, \langle H_J^{dc}(\tau) \rangle_0 + \frac{1}{2!} \int_{|\tau - \tau'| > \tau_c = D^{-1}} d\tau d\tau' \, \langle T \left[H_J^{dc}(\tau) H_J^{dc}(\tau') \right] \rangle_0 + \cdots \right\}, \tag{C2}$$

where we have made explicit the constraints on τ imposed by the finite bandwidth of the continuum of states described by Φ_{σ} and Φ_{σ}^{\dagger} .

Next, let us integrate out the high energy degrees of freedom contained $\Phi_{0\sigma}$ and $\Phi_{0\sigma}^{\dagger}$ (recall that $d_{\sigma}, d_{\sigma}^{\dagger}$ describe a low-energy subgap state and it cannot be integrated out). Such degrees of freedom involve excitations with energies $\sim D$ above the ground state and therefore determine the short imaginary time behavior of the Green's functions for Φ_{σ} . Note that, since at excitation energies $\sim D$ Bogoliubov quasi-particles either behave as electrons or holes (in other words, either $u \to 0$ or $v \to 0$), the anomalous GFs involving the operator Φ_0 , i.e. $\langle \mathcal{T} [\Phi_{0\uparrow}(\tau) \Phi_{0\downarrow}(\tau')] \rangle_0$, etc, vanish for $|\tau - \tau'| \simeq \tau_c^{-1}$. Thus, in the above perturbation series, for $|\tau - \tau'| \sim \tau_c$, we need to consider only normal correlations, which take the familiar Fermi liquid form:

$$\langle \mathcal{T} \left[\Phi_{0\sigma}(\tau) \Phi_{0\sigma'}^{\dagger}(\tau') \right] \rangle_0 \simeq \frac{\nu_0 \delta_{\sigma\sigma'}}{(\tau - \tau')}$$
 (C3)

for $|\tau' - \tau| \simeq \tau_c^{-1}$, where ν_0 is the (mean) density of states of the normal state. Thus, the first non-constant contribution to the scaling of the couplings stems from the second order term. We first split the integrals over τ, τ' according to:

$$\int_{\tau-\tau'|>\tau_c=D^{-1}} d\tau d\tau' \dots = \int_{|\tau-\tau'|>\tau'_c=(D')^{-1}} d\tau d\tau' \dots + \int_{\tau'_c=(D')^{-1}>|\tau-\tau'|>\tau_c=D^{-1}} d\tau d\tau' \dots$$
(C4)

and consider the terms in the second term for which $\tau'_c > |\tau - \tau'| > \tau_c$. Expanding the second order term in powers, corrections to the couplings contained in the first order term are generated at $O(J^2_{d\Phi})$, $O(J_{d\Phi}J_{\Phi\Phi})$ and $O(J^2_{\Phi\Phi})$. We explicitly evaluate below the $O(J^2_{d\Phi})$ term. The calculations for the remaining terms are similar and not reproduced

I

here. Einstein's convention of repeated index summation is used throughout:

$$O(J_{d\Phi}^{2}) = \frac{J_{d\Phi}^{2}(D)}{2!} \int_{\tau_{c}^{\prime} > |\tau - \tau^{\prime}| > \tau_{c}} d\tau d\tau^{\prime} \left\{ \langle \mathcal{T} \left[S^{a}(\tau) S^{b}(\tau^{\prime}) \right] \rangle_{0} \left(s_{\sigma\sigma^{\prime}}^{a} s_{\lambda\lambda^{\prime}}^{b} \right) \langle \mathcal{T} \left[d_{\sigma}^{\dagger}(\tau) \Phi_{0\sigma^{\prime}}(\tau) \Phi_{0\lambda}^{\dagger}(\tau^{\prime}) d_{\lambda^{\prime}}(\tau^{\prime}) \right] \rangle_{0} \right. \\ \left. + \left\langle \mathcal{T} \left[S^{a}(\tau) S^{b}(\tau^{\prime}) \right] \rangle_{0} \left(s_{\sigma\sigma^{\prime}}^{a} s_{\lambda\lambda^{\prime}}^{b} \right) \langle \mathcal{T} \left[\Phi_{0\sigma}^{\dagger}(\tau) d_{\sigma^{\prime}}(\tau) d_{\lambda}^{\dagger}(\tau^{\prime}) \Phi_{0\lambda^{\prime}}(\tau^{\prime}) \right] \rangle_{0} \right\}$$
(C5)

$$= \frac{J_{d\Phi}^{2}(D)\nu_{0}}{2!} \int_{\tau_{c}^{\prime} > |\tau-\tau^{\prime}| > \tau_{c}} d\tau d\tau^{\prime} \left\{ \langle \mathcal{T} \left[S^{a}(\tau)S^{b}(\tau^{\prime}) \right] \rangle_{0} \frac{\left(s_{\sigma\lambda}^{a} s_{\lambda\lambda^{\prime}}^{b} \right)}{(\tau-\tau^{\prime})} \langle \mathcal{T} \left[d_{\sigma}^{\dagger}(\tau) d_{\lambda^{\prime}}(\tau^{\prime}) \right] \rangle_{0} \right. \\ \left. + \left\langle \mathcal{T} \left[S^{a}(\tau)S^{b}(\tau^{\prime}) \right] \right\rangle_{0} \frac{\left(s_{\lambda\sigma}^{b} s_{\sigma\sigma^{\prime}}^{a} \right)}{(\tau-\tau^{\prime})} \langle \mathcal{T} \left[d_{\sigma^{\prime}}(\tau) d_{\lambda}^{\dagger}(\tau^{\prime}) \right] \rangle_{0} \right\}$$
(C6)

$$= -\frac{J_{d\Phi}^2(D)\nu_0}{4} \int d\tau d\tau' \frac{i\epsilon^{abc} \left[s^a, s^b\right]_{\sigma\sigma'}}{|\tau - \tau'|} \langle T \left[S^c(\tau)\right] \rangle_0 \langle \mathcal{T} \left[d_{\sigma}^{\dagger}(\tau)d_{\sigma'}(\tau')\right] \rangle_0 \tag{C7}$$

$$\tau_{c}^{\prime} > |\tau - \tau^{\prime}| > \tau_{c}$$

$$= -\frac{J_{d\Phi}^{2}(D)\nu_{0}}{2} \int d\tau d\tau^{\prime} \frac{\left(\epsilon^{abc}\epsilon^{abf}s_{\sigma\sigma^{\prime}}^{f}\right)}{|\tau - \tau^{\prime}|} \langle T\left[S^{c}(\tau)\right] \rangle_{0} \langle \mathcal{T}\left[d_{\sigma}^{\dagger}(\tau)d_{\sigma^{\prime}}(\tau^{\prime})\right] \rangle_{0} \tag{C8}$$

$$= -\frac{\sigma_{d\Phi}(D)\nu_{0}}{2} \int_{\tau_{c}' > |\tau - \tau'| > \tau_{c}} d\tau d\tau' \frac{\langle \tau \rangle \sigma \sigma'}{|\tau - \tau'|} \langle T[S^{c}(\tau)] \rangle_{0} \langle \mathcal{T}[d_{\sigma}^{\dagger}(\tau)d_{\sigma'}(\tau')] \rangle_{0}$$
(C8)

$$= -\frac{J_{d\Phi}^2(D)\nu_0}{2} \int_{\tau_c' > |\tau - \tau'| > \tau_c} d\tau d\tau' \frac{1}{|\tau - \tau'|} \langle \mathcal{T} \left[d_{\sigma}^{\dagger}(\tau) S^c(\tau) s_{\sigma\sigma'}^c d_{\sigma'}(\tau') \right] \rangle_0 \tag{C9}$$

In the above derivation we have used the following results: $\epsilon^{abc}\epsilon^{abd} = 2\delta_{cd}$ and

$$\mathcal{T}\left[S^{a}(\tau)S^{b}(\tau')\right] = \theta(\tau - \tau')S^{a}S^{b} + \theta(\tau' - \tau)S^{b}S^{a}$$
(C10)

$$= \frac{1}{2} (S^a S^b - S^b S^a) \left[\theta(\tau - \tau') - \theta(\tau' - \tau) \right] + \frac{1}{2} (S^a S^b + S^b S^a)$$
(C11)

$$=\frac{i}{2}\epsilon^{abc}S^{c}\operatorname{sgn}(\tau-\tau') + \left\{S^{a}, S^{b}\right\}$$
(C12)

because $S^a(\tau) = e^{H_0\tau}S^a e^{-H_0\tau} = S^a$. As noted above, the operators describing the dGSJ quasi-particle have time dynamics varying on the scale of $\Delta^{-1} \ll \tau'_c$, which is very slow compared to the fast degrees of freedom being integrated out from Φ_{0c} and Φ^{\dagger}_c . Introducing $\tau_- = \tau - \tau'$ and $\tau_+ = (\tau + \tau')/2$. Thus, the term proportional to $\{S^a, S^b\}$ drops because it is multiplied by τ_-^{-1} rather than $|\tau_-|^{-1}$ and the integral over τ_- of former vanishes to leading order. Thus, to leading order in τ_- , we are left with

$$O(J_{d\Phi}^2) = -J_{d\Phi}^2(D)\nu_0 \int d\tau_+ \left\langle \mathcal{T} \left[d_{\sigma}^{\dagger}(\tau_+) \mathbf{S}(\tau_+) \cdot \mathbf{s}_{\sigma\sigma'}(\tau_+) d_{\sigma}(\tau_+) \right] \right\rangle_0 \int_{\tau_c' > |\tau_-| > \tau_c} \frac{d\tau_-}{|\tau_-|} \tag{C13}$$

$$= -2\nu_0 \frac{\delta D}{D} J_{d\Phi}^2(D) \int d\tau \, \langle \mathcal{T} \left[d_{\sigma}^{\dagger}(\tau) \boldsymbol{S}(\tau) \cdot \boldsymbol{s}_{\sigma\sigma'}(\tau) d_{\sigma}(\tau) \right] \rangle_0. \tag{C14}$$

In the last expression, we have evaluated the integral over τ_{-} using

$$\int_{\tau_c' > |\tau_-| > \tau_c} \frac{d\tau_-}{|\tau_-|} = 2\log\left(\frac{\tau_c'}{\tau_c}\right) = 2\log\left(\frac{D}{D'}\right) = -2\log\left(\frac{D-\delta D}{D}\right) \simeq \frac{2\delta D}{D},\tag{C15}$$

and replaced $\tau_+ \to \tau$. Notice that the resulting expression in Eq. (C14) takes the same form as the contribution $\propto J_{dd}$ in the first order term of (C2). This leads to the following recursion relation:

$$J_{dd}(D - \delta D) = J_{dd}(D) + 2\nu_0 J_{d\Phi}^2(D) \frac{\delta D}{D}$$
(C16)

Assuming the couplings are continuous functions of the cut-off D, the recursion relation becomes a differential equation:

$$D\frac{dJ_{dd}(D)}{dD} = -2\nu_0 J_{d\Phi}^2,\tag{C18}$$

which implies that J_{cc} increases with decreasing bandwdith D.

Similarly, we can tackle the terms at $O(J_{d\Phi}J_{\Phi\Phi})$ and $O(J_{\Phi\Phi})$ (note the latter one is the only one present in the standard poor man's scaling treatment of the Kondo model). From those terms, the following differential equations are obtained:

$$D\frac{dJ_{d\Phi}(D)}{dD} = -2\nu_0 J_{d\Phi} J_{\Phi\Phi}, \tag{C19}$$

$$D\frac{dJ_{dd}(D)}{dD} = -2\nu_0 J_{\Phi\Phi}^2. \tag{C20}$$

It is convenient to introduce a new scaling variable defined by the differential equation:

$$\frac{dD}{D} = d\ell \Rightarrow D(\ell) = D_0 e^{-\ell}.$$
(C21)

Thus, as $\ell \to +\infty D(\ell) \to 0$. Furthermore, if we define the dimensionless couplings $g_{dd} = 2\nu_0 J_{dd}$, $g_{d\Phi} = 2\nu_0 J_{d\Phi}$, and $g_{\Phi\Phi} = 2\nu_0 J_{\Phi\Phi}$, we finally arrive at the scaling equations (14) to (16) discussed in Sec. V.

- ¹ L. Yu, Acta Physica Sinica **21**, 75 (1965).
- ² H. Shiba, Prog. Theor. Phys. **40**, 435 (1968).
- ³ A. I. Rusinov, Sov. J. Exp. Theor. Phys. **29**, 1101 (1969).
- ⁴ A. V. Balatsky, I. Vekhter, and J.-X. Zhu, Rev. Mod. Phys. **78**, 373 (2006).
- ⁵ S.-H. Ji, T. Zhang, Y.-S. Fu, X. Chen, X.-C. Ma, J. Li, W.-H. Duan, J.-F. Jia, and Q.-K. Xue, Phys. Rev. Lett. **100**, 226801 (2008).
- ⁶ B. W. Heinrich, J. I. Pascual, and K. J. Franke, Progress in Surface Science **93**, 1 (2018).
- ⁷ J. Kondo, Progress of Theoretical Physics **32**, 37 (1964).
- ⁸ A. Skurativska, J. Ortuzar, D. Bercioux, F. S. Bergeret, and M. A. Cazalilla, "Robust spin polarization of yushiba-rusinov states in superconductor/ferromagnetic insulator heterostructures," (2023), arXiv:2302.13864 [condmat.supr-con].
- ⁹ C. Rubio-Verdú, J. Zaldívar, R. Žitko, and J. I. Pascual, Phys. Rev. Lett. **126**, 017001 (2021).
- ¹⁰ K. Satori, H. Shiba, O. Sakai, and Y. Shimizu, Journal of the Physical Society of Japan **61**, 3239 (1992).
- ¹¹ R. žitko, O. Bodensiek, and T. Pruschke, Phys. Rev. B 83, 054512 (2011).
- ¹² A. Odobesko, D. Di Sante, A. Kowalski, S. Wilfert, F. Friedrich, R. Thomale, G. Sangiovanni, and M. Bode, Phys. Rev. B **102**, 174504 (2020).
- ¹³ I. Affleck, J.-S. Caux, and A. M. Zagoskin, Phys. Rev. B 62, 1433 (2000).
- ¹⁴ E. Vecino, A. Martín-Rodero, and A. L. Yeyati, Phys. Rev. B 68, 035105 (2003).
- ¹⁵ F. von Oppen and K. J. Franke, Phys. Rev. B **103**, 205424 (2021).
- ¹⁶ E. Liebhaber, L. M. Rütten, G. Reecht, J. F. Steiner, S. Rohlf, K. Rossnagel, F. von Oppen, and K. J. Franke, Nature Communications 13, 2160 (2022).
- ¹⁷ T. Machida, Y. Nagai, and T. Hanaguri, Phys. Rev. Res. 4, 033182 (2022).
- ¹⁸ S. Trivini, J. Ortuzar, K. Vaxevani, J. Li, F. S. Bergeret,

M. A. Cazalilla, and J. I. Pascual, Phys. Rev. Lett. **130**, 136004 (2023).

- ¹⁹ K. D. Usadel, Phys. Rev. Lett. **25**, 507 (1970).
- ²⁰ F. Zhou, P. Charlat, B. Spivak, and B. Pannetier, Journal of Low Temperature Physics **110**, 841 (1998).
- ²¹ H. le Sueur, P. Joyez, H. Pothier, C. Urbina, and D. Esteve, Phys. Rev. Lett. **100**, 197002 (2008).
- ²² A. K. Gupta, L. Crétinon, N. Moussy, B. Pannetier, and H. Courtois, Phys. Rev. B 69, 104514 (2004).
- ²³ P. Wei, S. Manna, M. Eich, P. Lee, and J. Moodera, Phys. Rev. Lett. **122**, 247002 (2019).
- ²⁴ L. Serrier-Garcia, J. C. Cuevas, T. Cren, C. Brun, V. Cherkez, F. Debontridder, D. Fokin, F. S. Bergeret, and D. Roditchev, Phys. Rev. Lett. **110**, 157003 (2013).
- ²⁵ K. Vaxevani, J. Li, S. Trivini, J. Ortuzar, D. Longo, D. Wang, and J. I. Pascual, Nano Lett. **22**, 6075 (2022).
- ²⁶ L. Schneider, K. T. Ton, I. Ioannidis, J. Neuhaus-Steinmetz, T. Posske, R. Wiesendanger, and J. Wiebe, "Proximity superconductivity in atom-by-atom crafted quantum dots," (2022).
- ²⁷ J.-C. Liu, R. Pawlak, X. Wang, P. D'Astolfo, C. Drechsel, P. Zhou, S. Decurtins, U. Aschauer, S.-X. Liu, W. Wulfhekel, and E. Meyer, "Proximity-induced superconductivity in atomically precise nanographene," (2022).
- ²⁸ Y. Zhao, K. Jiang, C. Li, Y. Liu, G. Zhu, M. Pizzochero, E. Kaxiras, D. Guan, Y. Li, H. Zheng, C. Liu, J. Jia, M. Qin, X. Zhuang, and S. Wang, Nature Chemistry (2022), 10.1038/s41557-022-01061-5.
- ²⁹ S. Mishra, G. Catarina, F. Wu, R. Ortiz, D. Jacob, K. Eimre, J. Ma, C. A. Pignedoli, X. Feng, P. Ruffieux, J. Fernández-Rossier, and R. Fasel, Nature **598**, 287 (2021).
- ³⁰ J. Hieulle, S. Castro, N. Friedrich, A. Vegliante, F. R. Lara, S. Sanz, D. Rey, M. Corso, T. Frederiksen, J. I. Pascual, and D. Peña, Angewandte Chemie International Edition **60**, 25224 (2021).
- ³¹ S. Mishra, D. Beyer, K. Eimre, R. Ortiz, J. Fernández-

Rossier, R. Berger, O. Gröning, C. A. Pignedoli, R. Fasel, X. Feng, and P. Ruffieux, Angewandte Chemie International Edition **59**, 12041 (2020).

- ³² P. G. de Gennes and D. Saint-James, Phys. Lett. 4 (1963), 10.1016/0031-9163(63)90148-3.
- ³³ I. O. Kulik, Soviet Journal of Experimental and Theoretical Physics **30**, 944 (1969).
- ³⁴ E. Cortés-del Río, J. L. Lado, V. Cherkez, P. Mallet, J.-Y. Veuillen, J. C. Cuevas, J. M. Gómez-Rodríguez, J. Fernández-Rossier, and I. Brihuega, Advanced Materials **33**, 2008113 (2021).
- ³⁵ J. O. Island, R. Gaudenzi, J. de Bruijckere, E. Burzurí, C. Franco, M. Mas-Torrent, C. Rovira, J. Veciana, T. M.

Klapwijk, R. Aguado, and H. S. J. van der Zant, Phys. Rev. Lett. **118**, 117001 (2017).

- ³⁶ P. W. Anderson, Journal of Physics C: Solid State Physics 3, 2436 (1970).
- ³⁷ P. G. DE GENNES, Rev. Mod. Phys. **36**, 225 (1964).
- ³⁸ W. L. McMillan, Phys. Rev. **175**, 559 (1968).
- ³⁹ G. Kieselmann, Physical Review B **35**, 6762 (1987).
- ⁴⁰ G. B. Arnold, Phys. Rev. B **18**, 1076 (1978).
- ⁴¹ G. Kieselmann, Phys. Rev. B **35**, 6762 (1987).
- ⁴² J. Rowell and W. McMillan, Physical Review Letters 16, 453 (1966).
- ⁴³ W. Tomasch, Physical Review Letters **16**, 16 (1966).
- ⁴⁴ T. E. Feuchtwang, Phys. Rev. B **10**, 4121 (1974).